

# The Experimental Ideal

EC 607, Set 02

---

Edward Rubin  
Spring 2021

# Prologue

# Schedule

## Last time

Research basics, our class, and R

## Today

**Admin:** Zoom recordings on Canvas.

**Material:** The Rubin causal model (not mine), [Chapter 2 MHE](#).

**Assignment<sub>1</sub>** Make sure R and RStudio are running on your computer.

**Assignment<sub>2</sub>** Take 15 minutes to quietly think about your interests.

## Future

**Lab:** Meet Philip and start deepening R knowledge.

**Long run:** Deepen understandings/intuitions for causality and inference.

# Review

*Research fundamentals*

# Review

## Research fundamentals

Angrist and Pischke provide four **fundamental questions for research**:

1. What is the **causal relationship of interest**?
2. How would an **ideal experiment** capture this causal effect of interest?
3. What is your **identification strategy**?
4. What is your **mode of inference**?

Seemingly straightforward questions can be fundamentally unanswerable.

# Review

## General research recommendations

More unsolicited advice:

- Be curious.
- Ask questions.
- Attend seminars.
- Meet faculty (UO + visitors).
- Focus on learning—especially intuition.<sup>†</sup>
- **Be kind and constructive.**

<sup>†</sup> *Learning* is not always the same as getting good grades.

# The experimental ideal

# The experimental ideal

## What's so great about experiments?

Science widely regards **experiments as the gold standard** for research.

*But why?* The costs can be substantial.

### **Costs**

- slow and expensive
- heavily regulated by (risk-averse?) review boards
- can abstract away from the actual question/setting

### **Benefits**

So the benefits need to be pretty large, right?



# The experimental ideal

## Example: Hospitals and health

Imagine we want to know the **causal effect of hospitals on health**.

### Research question

Within the population of poor, elderly individuals, does visiting the emergency room for primary care improve health?

### Empirical exercise

1. Collect data on *health status* and *hospital visits*.
2. Summarize health status by hospital-visit group.

# The experimental ideal

## Example: Hospitals and health

Our empirical exercise from the 2005 National Health Interview Survey:

Group	Sample Size	Mean Health Status	Std. Error
Hospital	7,774	3.21	0.014
No hospital	90,049	3.93	0.003

We get a  $t$  statistic of 58.9 when testing a difference in groups' means (0.72).

**Conclusion?** Hospitals make folks worse. Hospitals make sick people sicker.

**Alternative conclusion:** Perhaps we're making a mistake in our analysis... maybe sick people go to hospitals?

# The experimental ideal

## Potential outcomes framework

Let's develop a framework to better discuss the problem here.

- Binary treatment variable (*e.g.*, hospitalized):  $D_i = 0, 1$
- Outcome for individual  $i$  (*e.g.*, health):  $Y_i$

This framework has a few names...

- Neyman potential outcomes framework
- Rubin causal model
- Neyman-Rubin `"potential outcome"|"causal" "framework"|"model"`

# The experimental ideal

## Potential outcomes framework

**Research question:** Does  $D_i$  affect  $Y_i$ ?

For each individual  $i$ , there are two **potential outcomes** (w/ binary  $D_i$ )

1.  $Y_{1i}$  if  $D_i = 1$

$i$ 's health outcome if she went to the hospital

2.  $Y_{0i}$  if  $D_i = 0$

$i$ 's health outcome if she did not go to the hospital

The difference between these two outcomes gives us the **causal effect of hospital treatment**, *i.e.*,

$$\tau_i = Y_{1i} - Y_{0i}$$

# The experimental ideal

## #problems

This simple equation

$$\tau_i = Y_{1i} - Y_{0i}$$

leads us to ***the fundamental problem of causal inference.***

■ We can never simultaneously observe  $Y_{1i}$  and  $Y_{0i}$ .

Most of applied econometrics focuses on addressing this simple problem.

Accordingly, our methods try to address the related question

■ For each  $Y_{1i}$ , what is a (reasonably) good counterfactual?

# The experimental ideal

## Solutions?

**Problem** We cannot directly calculate  $\tau_i = Y_{1i} - Y_{0i}$ .

### Proposed solution

Compare **outcomes for people who visited the hospital** ( $Y_{1i} \mid D_i = 1$ )  
to **outcomes for people who did not visit the hospital** ( $Y_{0j} \mid D_j = 0$ ).

$$E[Y_i \mid D_i = 1] - E[Y_i \mid D_i = 0]$$

which gives us the *observed difference in health outcomes*.

**Q** This comparison will return *an* answer, but is it *the* answer we want?

# The experimental ideal

## Selection

**Q** What does  $E[Y_i | D_i = 1] - E[Y_i | D_i = 0]$  actually tell us?

**A** First notice that we can write  $i$ 's outcome  $Y_i$  as

$$Y_i = Y_{0i} + D_i \underbrace{(Y_{1i} - Y_{0i})}_{\tau_i}$$

Now write out our expectation, apply this definition, do creative math.

$$\begin{aligned} & E[Y_i | D_i = 1] - E[Y_i | D_i = 0] \\ &= E[Y_{1i} | D_i = 1] - E[Y_{0i} | D_i = 0] \\ &= \underbrace{E[Y_{1i} | D_i = 1] - E[Y_{0i} | D_i = 1]}_{\text{Average treatment effect on the treated 😊}} + \underbrace{E[Y_{0i} | D_i = 1] - E[Y_{0i} | D_i = 0]}_{\text{Selection bias 😞}} \end{aligned}$$

# The experimental ideal

## Selection

The **first term** is *good variation*—essentially the answer that we want.

$$\begin{aligned} & E[Y_{1i} \mid D_i = 1] - E[Y_{0i} \mid D_i = 1] \\ &= E[Y_{1i} - Y_{0i} \mid D_i = 1] \\ &= E[\tau_i \mid D_i = 1] \end{aligned}$$

The **average causal effect** of hospitalization *for hospitalized individuals*.

The **second term** is bad variation—preventing us from knowing the answer.

$$E[Y_{0i} \mid D_i = 1] - E[Y_{0i} \mid D_i = 0]$$

The difference in the average untreated outcome between the treatment and control groups.

**Selection bias** The extent to which the "control group" provides a bad counterfactual for the treated individuals.



# The experimental ideal

## Selection

Angrist and Pischke (MHE, p. 15),

The goal of most empirical economic research is to overcome selection bias, and therefore to say something about the causal effect of a variable like  $\mathbf{D}_i$ .

**Q** So how do experiments—the gold standard of empirical economic (and scientific) research—accomplish this goal and overcome selection bias?

# The experimental ideal

## Back to experiments

**Q** How do experiments overcome selection bias?

**A** Experiments break the link between potential outcomes and treatment.

*In other words:* Randomly assigning  $\mathbf{D}_i$  makes  $\mathbf{D}_i$  independent of which outcome we observe (meaning  $\mathbf{Y}_{1i}$  or  $\mathbf{Y}_{0i}$ ).

**Difference in means** with random assignment of  $\mathbf{D}_i$

$$\begin{aligned} & E[\mathbf{Y}_i \mid \mathbf{D}_i = 1] - E[\mathbf{Y}_i \mid \mathbf{D}_i = 0] \\ &= E[\mathbf{Y}_{1i} \mid \mathbf{D}_i = 1] - E[\mathbf{Y}_{0i} \mid \mathbf{D}_i = 0] \\ &= E[\mathbf{Y}_{1i} \mid \mathbf{D}_i = 1] - E[\mathbf{Y}_{0i} \mid \mathbf{D}_i = 1] \quad \text{from random assignment of } \mathbf{D}_i \\ &= E[\mathbf{Y}_{1i} - \mathbf{Y}_{0i} \mid \mathbf{D}_i = 1] \\ &= E[\boldsymbol{\tau}_i \mid \mathbf{D}_i = 1] \\ &= E[\boldsymbol{\tau}_i] \quad \text{Random assignment of } \mathbf{D}_i \text{ breaks selection bias.} \end{aligned}$$

# The experimental ideal

## Randomly assigned treatment

The key to avoiding selection bias: **random assignment of treatment** (or *as-good-as random assignment*, e.g., natural experiments).

Random assignment of treatment gives us

$$E[Y_{0i} | D_i = 0] = E[Y_{0i} | D_i = 1]$$

meaning the control group's mean now provides a good counterfactual for the treatment group's mean.

In other words, there is no selection bias, *i.e.*,

$$\text{Selection bias} = E[Y_{0i} | D_i = 1] - E[Y_{0i} | D_i = 0] = 0$$

# The experimental ideal

## Randomly assigned treatment

Additional benefit of randomization:

The *average treatment effect* is now representative of the *population average*, rather than the *treatment-group average*.

$$E[\tau_i \mid \mathbf{D}_i = 1] = E[\tau_i \mid \mathbf{D}_i = 0] = E[\tau_i]$$

# The experimental ideal

## Example: Training programs

Governments subsidize training programs to assist disadvantaged workers.

**Q** Do these programs have the desired effects (*i.e.*, increase wages)?

**A** Observational studies—comparing wage data from participants and non-participants—often find that people who complete these programs actually make **lower wages**.

**Challenges** Participants self select. + Programs target lower-wage workers.

# The experimental ideal

## Example: Training programs

How do we formalize these concerns in our framework?

### Observational program evaluations

$$\begin{aligned} E[\text{Wage}_i \mid \text{Program}_i = 1] - E[\text{Wage}_i \mid \text{Program}_i = 0] = \\ \underbrace{E[\text{Wage}_{1i} \mid \text{Program}_i = 1] - E[\text{Wage}_{0i} \mid \text{Program}_i = 1]}_{\text{Average causal effect of training program on wages for participants, i.e., } \bar{\tau}_1} + \\ \underbrace{E[\text{Wage}_{0i} \mid \text{Program}_i = 1] - E[\text{Wage}_{0i} \mid \text{Program}_i = 0]}_{\text{Selection bias}} \end{aligned}$$

If the program attracts/selects individuals who, on average, have lower wages without the program (sort of the point of the program), then we have negative selection bias.

# The experimental ideal

## Example: Training programs

$$\begin{aligned} E[\text{Wage}_i \mid \text{Program}_i = 1] - E[\text{Wage}_i \mid \text{Program}_i = 0] = \\ E[\text{Wage}_{1i} \mid \text{Program}_i = 1] - E[\text{Wage}_{0i} \mid \text{Program}_i = 1] + \\ E[\text{Wage}_{0i} \mid \text{Program}_i = 1] - E[\text{Wage}_{0i} \mid \text{Program}_i = 0] \end{aligned}$$

So even if the program, on average, has a positive wage effect (in the participant group), *i.e.*,  $\bar{\tau}_1 > 0$ , we will detect a lower effect due to the negative selection bias.

If the bias is sufficiently large (relative to the treatment effect), our estimate will even get the sign of the effect wrong.

**Related** While observational studies typically found negative program effects, several experiments found positive program effects.

# The experimental ideal

## Example: The STAR experiment

The Tennessee STAR experiment is a famous/popular example of an experiment that allows us to answer an important social/policy question.

**Research question** Do classroom resources affect student performance?

- Statewide(-ish) in Tennessee for the 1985–1986 kindergarten cohort
- Ran for 4 years with ~11,600 children. Cost ~\$12 million.

### Treatments

1. *Small* classes (13–17 students)
2. *Regular* classes (22–35 students) plus part-time teacher's aide
3. *Regular* classes (22–35 students) plus full-time teacher's aide



# The experimental ideal

## Example: The STAR experiment

**First question** Did the randomization balance participants' characteristics across the treatment groups?

Ideally, we would have pre-experiment data on outcome variable.

Unfortunately, we only have a few demographic attributes.

Table 2.2.1, MHE

<b>Treatment: Class Size</b>				
<b>Variable</b>	<b>Small</b>	<b>Regular</b>	<b>Regular + Aide</b>	<b>P-value</b>
<i>Free lunch</i>	0.47	0.48	0.50	0.09
<i>White/Asian</i>	0.68	0.67	0.66	0.26
<i>Age in 1985</i>	5.44	5.43	5.42	0.32
<i>Attrition rate</i>	0.49	0.52	0.53	0.02
<i>K. class size</i>	15.10	22.40	22.80	0.00
<i>K. test percentile</i>	54.70	48.90	50.00	0.00

Table 2.2.1, MHE

<b>Variable</b>	<b>Treatment: Class Size</b>			<b>P-value</b>
	<b>Small</b>	<b>Regular</b>	<b>Regular + Aide</b>	
<i><b>Free lunch</b></i>	<b>0.47</b>	<b>0.48</b>	<b>0.50</b>	<b>0.09</b>
<i><b>White/Asian</b></i>	<b>0.68</b>	<b>0.67</b>	<b>0.66</b>	<b>0.26</b>
<i><b>Age in 1985</b></i>	<b>5.44</b>	<b>5.43</b>	<b>5.42</b>	<b>0.32</b>
<i>Attrition rate</i>	0.49	0.52	0.53	0.02
<i>K. class size</i>	15.10	22.40	22.80	0.00
<i>K. test percentile</i>	54.70	48.90	50.00	0.00

Demographics appear balanced across the three treatment groups.

Table 2.2.1, MHE

<b>Variable</b>	<b>Treatment: Class Size</b>			<b>P-value</b>
	<b>Small</b>	<b>Regular</b>	<b>Regular + Aide</b>	
<i>Free lunch</i>	0.47	0.48	0.50	0.09
<i>White/Asian</i>	0.68	0.67	0.66	0.26
<i>Age in 1985</i>	5.44	5.43	5.42	0.32
<b><i>Attrition rate</i></b>	<b>0.49</b>	<b>0.52</b>	<b>0.53</b>	<b>0.02</b>
<i>K. class size</i>	15.10	22.40	22.80	0.00
<i>K. test percentile</i>	54.70	48.90	50.00	0.00

The three groups differ significantly on attrition rate.

Table 2.2.1, MHE

<b>Variable</b>	<b>Treatment: Class Size</b>			<b>P-value</b>
	<b>Small</b>	<b>Regular</b>	<b>Regular + Aide</b>	
<i>Free lunch</i>	0.47	0.48	0.50	0.09
<i>White/Asian</i>	0.68	0.67	0.66	0.26
<i>Age in 1985</i>	5.44	5.43	5.42	0.32
<i>Attrition rate</i>	0.49	0.52	0.53	0.02
<b><i>K. class size</i></b>	<b>15.10</b>	<b>22.40</b>	<b>22.80</b>	<b>0.00</b>
<i>K. test percentile</i>	54.70	48.90	50.00	0.00

The randomization generated variation in the treatment.

Table 2.2.1, MHE

<b>Variable</b>	<b>Treatment: Class Size</b>			<b>P-value</b>
	<b>Small</b>	<b>Regular</b>	<b>Regular + Aide</b>	
<i>Free lunch</i>	0.47	0.48	0.50	0.09
<i>White/Asian</i>	0.68	0.67	0.66	0.26
<i>Age in 1985</i>	5.44	5.43	5.42	0.32
<i>Attrition rate</i>	0.49	0.52	0.53	0.02
<i>K. class size</i>	15.10	22.40	22.80	0.00
<b><i>K. test percentile</i></b>	<b>54.70</b>	<b>48.90</b>	<b>50.00</b>	<b>0.00</b>

The small-class treatment significantly increased test scores.

# The experimental ideal

## The STAR experiment

The previous table estimated/compared the treatment effects using simple differences in means.

We can make the same comparisons using regressions.

Specifically, we regress our outcome (test percentile) on dummy variables (binary indicator variables) for each treatment group.

# The experimental ideal

Example of our three treatment dummies.

$i$	$y_i$	$\text{Trt}_{1i}$	$\text{Trt}_{2i}$	$\text{Trt}_{3i}$
1	$y_1$	1	0	0
2	$y_2$	1	0	0
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$\ell$	$y_\ell$	1	0	0
$\ell + 1$	$y_{\ell-1}$	0	1	0
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$p$	$y_p$	0	1	0
$p + 1$	$y_{p+1}$	0	0	1
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$N$	$y_N$	0	0	1



# The experimental ideal

## Regression analysis

Assume for the moment that the treatment effect is constant<sup>†</sup>, *i.e.*,

$$Y_{1i} - Y_{0i} = \rho \quad \forall i$$

then we can rewrite

$$Y_i = Y_{0i} + D_i (Y_{1i} - Y_{0i})$$

as

$$Y_i = \underbrace{\alpha}_{=E[Y_{0i}]} + D_i \underbrace{\rho}_{Y_{1i} - Y_{0i}} + \underbrace{\eta_i}_{Y_{0i} - E[Y_{0i}]}$$

<sup>†</sup>You'll often hear econometricians say "homogeneous" (vs. "heterogeneous").

# The experimental ideal

## Regression analysis

$$Y_i = \alpha + D_i\rho + \eta_i$$

Now write out the conditional expectation of  $Y_i$  for both levels of  $D_i$

$$E[Y_i \mid D_i = 1] = E[\alpha + \rho + \eta_i \mid D_i = 1] = \alpha + \rho + E[\eta_i \mid D_i = 1]$$

$$E[Y_i \mid D_i = 0] = E[\alpha + \eta_i \mid D_i = 0] = \alpha + E[\eta_i \mid D_i = 0]$$

Take the difference...

$$\begin{aligned} E[Y_i \mid D_i = 1] - E[Y_i \mid D_i = 0] \\ = \rho + \underbrace{E[\eta_i \mid D_i = 1] - E[\eta_i \mid D_i = 0]}_{\text{Selection bias}} \end{aligned}$$

# The experimental ideal

## Regression analysis

$$E[Y_i | \mathbf{D}_i = 1] - E[Y_i | \mathbf{D}_i = 0] = \rho + E[\eta_i | \mathbf{D}_i = 1] - E[\eta_i | \mathbf{D}_i = 0]$$

Again, our estimate of the **treatment effect** ( $\rho$ ) is only going to be as good as our ability to shut down the **selection bias**.

**Selection bias in regression model:**  $E[\eta_i | \mathbf{D}_i = 1] - E[\eta_i | \mathbf{D}_i = 0]$

Selection bias here should remind you a lot of **omitted-variable bias**.

There is something in our disturbance  $\eta_i$  that is affecting  $\mathbf{Y}_i$  and is also correlated with  $\mathbf{D}_i$ .

In other metrics-y words: Our treatment  $\mathbf{D}_i$  is endogenous.

# The experimental ideal

## Solutions and covariates

**Selection bias in regression model:**  $E[\eta_i | \mathbf{D}_i = 1] - E[\eta_i | \mathbf{D}_i = 0]$

As before, if we randomly assign  $\mathbf{D}_i$ , then selection bias disappears.

Another potential route to identification is to condition on covariates in the hopes that they "take care of" the relationship between  $\mathbf{D}_i$  and whatever is in our disturbance  $\eta_i$ .

Without very clear reasons explaining how you know you've controlled for the "bad variation", clean and convincing identification on this path is going to be challenging.

# The experimental ideal

## Covariates

That said, covariates can help with two things:

1. Even experiments may need **conditioning/controls**: The STAR experiment was random *within school*—not across schools.
2. Covariates can soak up unexplained variation—**increasing precision**.

Now that we've seen regression can analyze experiments, let's estimate the STAR example...

Table 2.2.2, MHE

<b>Explanatory variable</b>	<b>1</b>	<b>2</b>	<b>3</b>
<i>Small class</i>	4.82 (2.19)	5.37 (1.26)	5.36 (1.21)
<i>Regular + aide</i>	0.12 (2.23)	0.29 (1.13)	0.53 (1.09)
<i>White/Asian</i>			8.35 (1.35)
<i>Female</i>			4.48 (0.63)
<i>Free lunch</i>			-13.15 (0.77)
<i>School F.E.</i>	F	T	T

The omitted level is *Regular* (with part-time aide).

Table 2.2.2, MHE

<b>Explanatory variable</b>	<b>1</b>	<b>2</b>	<b>3</b>
<i>Small class</i>	<b>4.82</b> (2.19)	5.37 (1.26)	5.36 (1.21)
<i>Regular + aide</i>	<b>0.12</b> (2.23)	0.29 (1.13)	0.53 (1.09)
<i>White/Asian</i>			8.35 (1.35)
<i>Female</i>			4.48 (0.63)
<i>Free lunch</i>			-13.15 (0.77)
<i>School F.E.</i>	<b>F</b>	T	T

Results without other controls are very similar to the difference in means.

Table 2.2.2, MHE

<b>Explanatory variable</b>	<b>1</b>	<b>2</b>	<b>3</b>
<i>Small class</i>	4.82 (2.19)	<b>5.37</b> <b>(1.26)</b>	5.36 (1.21)
<i>Regular + aide</i>	0.12 (2.23)	<b>0.29</b> <b>(1.13)</b>	0.53 (1.09)
<i>White/Asian</i>			8.35 (1.35)
<i>Female</i>			4.48 (0.63)
<i>Free lunch</i>			-13.15 (0.77)
<i>School F.E.</i>	F	<b>T</b>	T

School FEs enforce the experiment's design and increase precision.



Table 2.2.2, MHE

<b>Explanatory variable</b>	<b>1</b>	<b>2</b>	<b>3</b>
<i>Small class</i>	4.82 (2.19)	5.37 (1.26)	<b>5.36</b> <b>(1.21)</b>
<i>Regular + aide</i>	0.12 (2.23)	0.29 (1.13)	<b>0.53</b> <b>(1.09)</b>
<i>White/Asian</i>			<b>8.35</b> <b>(1.35)</b>
<i>Female</i>			<b>4.48</b> <b>(0.63)</b>
<i>Free lunch</i>			<b>-13.15</b> <b>(0.77)</b>
<i>School F.E.</i>	F	T	<b>T</b>

Additional controls slightly increase precision.

# Table of contents

## Admin

1. Schedule
2. Review

## Experimental ideal

1. Neyman/Rubin framework
2. Selection
3. Experiments
4. Example: Training programs
5. Example: STAR experiment
  - Mean differences
  - Dummy variables
  - Regression analysis
  - Covariates